

Quantum simulation of quantum chemistry

A case study using IBM Q

Carlos Tavares Sofia Oliveira Vitor Fernandes Mikhail Vasilevsky

High-Assurance Software Laboratory/INESC TEC and Departamento de Física
Universidade do Minho

April 14, 2019
Q-Days, Universidade do Minho

Outline

- 1 Quantum Simulation
- 2 Modeling Quantum Systems
 - Hamiltonian Formalism
 - Second Quantization
- 3 Building Quantum Circuits
 - Jordan-Wigner Transformation
 - Quantum Circuit and Tools
- 4 Techniques
 - Variational Quantum Eigensolver
- 5 Experiments
 - IBM
 - Results
 - H₂ molecule
 - LiH molecule
- 6 Expectations for the Future

Motivation

Simulation of quantum systems is computationally hard!

Motivation

Quantum Simulation was idealized by Feynman¹ and refined by Lloyd².

Quote of Richard Feynman:

“Nature isn’t classical, dammit, and if you want to make a simulation of nature, you’d better make it quantum mechanical, and by golly, it’s a wonderful problem because it doesn’t look so easy.”

¹[Richard P Feynman. Simulating physics with computers. International journal of theoretical physics, 21(6-7):467–488, 1982.]

²[Seth Lloyd. Universal quantum simulators. Science, pages 1073–1078, 1996.]

Motivation

There are many applications for quantum Simulation¹:

- (...)
- Quantum chemistry
 - Open quantum systems
 - Condensed Matter physics
 - Nuclear physics, High-energy physics (Particle physics)
 - Cosmology!

Multiple strategies, from "digital" quantum simulations to quantum *analogues*!

¹[GEORGESCU, Iulia M.; ASHHAB, Sahel; NORI, Franco. Quantum simulation. Reviews of Modern Physics, 2014, 86.1: 153.]

Motivation



Figure: Picture of a black hole

[Alsing, Paul M., Jonathan P. Dowling, and G. J. Milburn. "Ion trap simulations of quantum fields in an expanding universe." Physical review letters 94.22 (2005): 220401.]

The Schrödinger equation

The Schrödinger equation

$$i\hbar \frac{d}{dt} \Psi = H\Psi$$

Classical physics

$$K = \frac{1}{2}mv^2$$

Quantum physics

$$K = \frac{\hbar^2}{2m} \frac{d^2}{dx^2}$$

In position basis:

$$i\hbar \frac{d}{dt} \Psi = \left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) \right) \Psi$$

Solutions for a free particle with $V(x) = 0$, with p being the momentum and E the energy of the system.

$$\Psi(x, t) = Ae^{-i(px - Et)/\hbar}$$

The hydrogen atom

Considering an *Hydrogen* atom (composed by 1 proton and 1 electron):

Figure: The hydrogen atom

Solutions of the Schrödinger equation¹ (3-dimensions):

$$\psi_{nlm}(r, \vartheta, \varphi) = \sqrt{\left(\frac{2}{na_0}\right) \frac{n-l-1!}{2n(n+l)!}} e^{-\frac{r}{a_0}} L_{n-l-1}^{2l+1}(p) Y_l^m(\vartheta, \varphi) \quad (1)$$

To every combination of n, l, m , denominated **quantum numbers**, corresponds a different solution of the equation.

¹[Y_l^m are the *spherical harmonics* functions and the L_{n-l-1}^{2l+1} are *Laguerre polynomials*]

The Schrödinger equation - interpretation

Which of those equations capture the behaviour of a system? Answer: A superposition of them?

- A quantum number refers to a physical characteristic of the system: energy, momentum, braiding...
- One (of several) problems: the **Measurement problem!**

Quantum mechanics is captured in the Hilbert space formalism:

- States $|\Psi\rangle$
 - ▶ States are vectors in the Hilbert space, spanned by an orthogonal basis $\langle\Psi_1|\Psi_2\rangle = 0$:

$$|\Psi\rangle = \lambda_1 |\Psi_1\rangle + \lambda_2 |\Psi_2\rangle + \dots + \lambda_n |\Psi_n\rangle$$

- ▶ **Superposition principle:** every superposition of the base vectors is a possible state of the system.

The Schrödinger equation - interpretation

- ▶ **Börn interpretation:** Probabilistic interpretation! Probability of obtaining $|\Psi_x\rangle$ given by $|\lambda_x|^2$ and $\sum_x |\lambda|^2 = 1$
- Operators $O|\Psi\rangle$
 - ▶ Operators are linear operators/matrix on the Hilbert space.

$$\begin{bmatrix} \langle \Psi_1 | O \Psi_1 \rangle & \langle \Psi_2 | O \Psi_1 \rangle & \dots & \langle \Psi_n | O \Psi_1 \rangle \\ \langle \Psi_1 | O \Psi_2 \rangle & \langle \Psi_2 | O \Psi_2 \rangle & \dots & \langle \Psi_n | O \Psi_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle \Psi_1 | O \Psi_n \rangle & \langle \Psi_2 | O \Psi_n \rangle & \dots & \langle \Psi_n | O \Psi_n \rangle \end{bmatrix} \quad (2)$$

What is quantum simulation ?

Objective:

- Mimic the **Hamiltonian operator**: Find an \hat{H} such that $H|\Psi\rangle \approx \hat{H}|\Psi\rangle$
- Mimic the **evolution of the Hamiltonian operator**: Find $e^{i\hat{H}}$ such that $e^{i\hat{H}}|\Psi(0)\rangle \approx e^{iH}|\Psi(0)\rangle!$

Evolution of a quantum operator (e^{iH} is a unitary operator)

$$|\Psi(t)\rangle = e^{iH}|\Psi(0)\rangle$$

Another way of looking into this (application of the operator over himself until the infinity):

$$e^{iH} = \lim_{n \rightarrow \infty} \left(I + \frac{iH}{n} \right)^n$$

Approximation methods: *Taylor* expansion, *Trotter* method

Computation

We are usually interested in calculate a properties of interest given Hamiltonians: lowest energy level of an Hamiltonian or other properties of the energy spectra, energy evolution throughout time ...

Finding the Ground State is hard even for a quantum computer, and in fact many classical optimization problems (NP-HARD) can be reduced to a ground state problem

The complexity of these problems are studied in the field of **Hamiltonian complexity**

- Class QMA is the quantum analogous of the NP class
- Problems: QC-SAT, QC-Circuit, k-Local Hamiltonian, Density matrix consistency

Hamiltonian Formalism

The hamiltonians describe the total energy of systems, which contains the operations associated with the kinetic and potential energies.

Generic formula:

$$H = K + V = -\frac{\hbar}{2m} \nabla^2 + V$$

For the majority of chemistry systems a Fermionic¹ Hamiltonian is enough. Total Fermionic Hamiltonian considering all interactions:

$$H = -\sum_{i=1}^N \frac{1}{2} \nabla_i^2 - \sum_{A=1}^M \frac{1}{2M_A} \nabla_{A^2} - \sum_{i=1}^N \sum_{A=1}^M \frac{Z_A}{r_{iA}} + \sum_{i=1}^N \sum_{j>i}^N \frac{1}{r_{ij}} + \sum_{A=1}^M \sum_{B>A}^M \frac{Z_A Z_B}{r_{BA}}$$

¹Fermions are subatomic particles, such as an electron/proton, which has half-integral spin and follows the Fermi-Dirac statistics.

² Components (left to right) - kinetic energy electrons; kinetic energy of nuclei; electric attraction nuclei - electrons; electric repulsion electrons; electric repulsion nuclei

[Attila Szabo and Neil S Ostlund. Modern quantum chemistry: introduction to advanced electronic structure theory. Courier Corporation, 2012.]

Hamiltonian Formalism

The hamiltonians describe the total energy of systems, which contains the operations associated with the kinetic and potential energies.

Generic formula:

$$H = K + V = -\frac{\hbar}{2m}\nabla^2 + V$$

For the majority of chemistry systems a Fermionic¹ Hamiltonian is enough. Electronic hamiltonian after applying Born-Oppenheimer approximation:

$$H_{elec} = -\sum_{i=1}^N \frac{1}{2} \nabla_i^2 - \sum_{i=1}^N \sum_{A=1}^M \frac{Z_A}{r_{iA}} + \sum_{i=1}^N \sum_{j>i}^N \frac{1}{r_{ij}}$$

¹Fermions are subatomic particles, such as an electron/proton, which has half-integral spin and follows the Fermi-Dirac statistics.

² Components (left to right) - kinetic energy electrons; kinetic energy of nuclei; electric attraction nuclei - electrons; electric repulsion electrons; electric repulsion nuclei

[Attila Szabo and Neil S Ostlund. Modern quantum chemistry: introduction to advanced electronic structure theory. Courier Corporation, 2012.]

Second Quantization

When dealing with Fermionic systems, *spin* plays an important role (Pauli exclusion principle):

$$\Psi(\chi_1, \chi_2) = -\Psi(\chi_2, \chi_1) \quad (3)$$

χ - Orbital, solution for the wave equation + spin, for a particle in the system

The Second Quantization is an alternative formalism for specifying Hamiltonians.

- Creation and annihilation operators (allows systems with variable dimension)

$$a_i^\dagger |\chi_1 \chi_2 \dots \chi_n\rangle = |\chi_i \chi_1 \chi_2 \dots \chi_n\rangle; a_i |\chi_i \chi_1 \chi_2 \dots \chi_n\rangle = |\chi_1 \chi_2 \dots \chi_n\rangle \quad (4)$$

[Dirac, Paul Adrien Maurice. The principles of quantum mechanics. No. 27. Oxford university press, 1981]

Second Quantization

Hamiltonian:

$$H = H_1 + H_2 = \sum_{\alpha, \beta=0}^{M-1} \tau_{\alpha\beta} a_{\alpha}^{\dagger} a_{\beta} + \frac{1}{2} \sum_{\alpha, \beta, \gamma, \delta=0}^{M-1} \mu_{\alpha\beta\gamma\delta} a_{\alpha}^{\dagger} a_{\gamma}^{\dagger} a_{\delta} a_{\beta}$$

where

$$\tau_{\alpha\beta} = \int dx_1 \psi_{\alpha}^*(x_1) \left(\frac{-\nabla^2}{2} + \sum_i \frac{Z_i}{|r_{i1}|} \right) \psi_{\beta}(x_1)$$

and

$$\mu_{\alpha\beta\gamma\delta} = \int dx_1 dx_2 \psi_{\alpha}^*(x_1) \psi_{\beta}(x_1) \left(\frac{1}{|r_{12}|} \right) \psi_{\gamma}^*(x_2) \psi_{\delta}(x_2)$$

where $\tau_{\alpha\beta}$ and $\mu_{\alpha\beta\gamma\delta}$ are matrix/operator coefficients.

What are the ψ s..?

- In our case, it has used STO-3G wavefunctions (ψ).

Jordan-Wigner Transformation

Has the purpose to map fermions into qubits.

σ^- represents the spin-lowering operator and σ^+ the spin-raising operator, which can be written in terms of Pauli operator:

$$\sigma^- = \frac{1}{2}(\sigma_x + i\sigma_y) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

$$\sigma^+ = \frac{1}{2}(\sigma_x - i\sigma_y) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Lowering and raising operators over sets of qubits:

$$a_j^\dagger = 1^{\otimes j-1} \otimes \sigma^- \otimes \sigma^{Z \otimes N-j-1}$$

$$a_j = 1^{\otimes j-1} \otimes \sigma^+ \otimes \sigma^{Z \otimes N-j-1}$$

[James D Whitfield, Jacob Biamonte, and Alán Aspuru-Guzik. Simulation of electronic structure hamiltonians using quantum computers. *Molecular Physics*, 109(5):735–750, 2011.]

Quantum Circuit

| Description | Second Quantization ^a | Pauli representation |
|--|---|--|
| Number Operator | $h_{pp} a_p^\dagger a_p$ | $\frac{h_{pp}}{2} (\mathbf{1}_p - \sigma_p^z)$ |
| Excitation Operator | $h_{pq} a_p^\dagger a_q + h_{qp} a_q^\dagger a_p$ | $\frac{1}{2} \left(\bigotimes_{k=q+1}^{p-1} \sigma_k^z \right) \left(\Re\{h_{pq}\} (\sigma_p^x \sigma_q^x + \sigma_p^y \sigma_q^y) + \Im\{h_{pq}\} (\sigma_p^y \sigma_q^x - \sigma_p^x \sigma_q^y) \right)$ |
| Coulomb Operators | $h_{pqqr} a_p^\dagger a_q^\dagger a_q a_p$ | $\frac{h_{pqqr}}{4} (\mathbf{1} - \sigma_p^z - \sigma_q^z + \sigma_p^z \sigma_q^z)$ |
| Number with ^b Excitation Operator | $h_{pqqr} a_p^\dagger a_q^\dagger a_q a_r + h_{rqpq} a_r^\dagger a_q^\dagger a_q a_p$ | $\left(\bigotimes_{k=r+1}^{p-1} \sigma_k^z \right) \left[\left(\Re\{h_{pqqr}\} (\sigma_r^x \sigma_p^x + \sigma_r^y \sigma_p^y) + \Im\{h_{pqqr}\} (\sigma_r^y \sigma_p^x - \sigma_r^x \sigma_p^y) \right) - \sigma_q^z \left(\Re\{h_{pqqr}\} (\sigma_r^x \sigma_p^x + \sigma_r^y \sigma_p^y) + \Im\{h_{pqqr}\} (\sigma_r^y \sigma_p^x - \sigma_r^x \sigma_p^y) \right) \right]$ |
| Double Excitation Operator | $h_{pqrs} a_p^\dagger a_q^\dagger a_r a_s + h_{srqp} a_s^\dagger a_r^\dagger a_q a_p$ | $\left(\bigotimes_{k=s+1}^{r-1} \sigma_k^z \right) \left(\bigotimes_{k=q+1}^{p-1} \sigma_k^z \right) \left(\frac{\Re\{h_{pqrs}\}}{8} \left(\begin{array}{l} \sigma_s^x \sigma_r^x \sigma_q^x \sigma_p^x - \sigma_s^x \sigma_r^x \sigma_q^y \sigma_p^y + \sigma_s^x \sigma_r^y \sigma_q^x \sigma_p^y \\ + \sigma_s^y \sigma_r^x \sigma_q^x \sigma_p^y + \sigma_s^y \sigma_r^x \sigma_q^y \sigma_p^x - \sigma_s^y \sigma_r^y \sigma_q^x \sigma_p^x \\ + \sigma_s^x \sigma_r^y \sigma_q^x \sigma_p^x + \sigma_s^y \sigma_r^y \sigma_q^y \sigma_p^y \end{array} \right) + \frac{\Im\{h_{pqrs}\}}{8} \left(\begin{array}{l} \sigma_s^y \sigma_r^x \sigma_q^x \sigma_p^x + \sigma_s^x \sigma_r^y \sigma_q^x \sigma_p^x - \sigma_s^x \sigma_r^x \sigma_q^y \sigma_p^y \\ - \sigma_s^x \sigma_r^y \sigma_q^y \sigma_p^y - \sigma_s^y \sigma_r^x \sigma_q^y \sigma_p^x + \sigma_s^y \sigma_r^y \sigma_q^x \sigma_p^y \\ + \sigma_s^y \sigma_r^y \sigma_q^x \sigma_p^x + \sigma_s^x \sigma_r^x \sigma_q^y \sigma_p^x \end{array} \right) \right)$ |

Figure: Summary of operators representation

In fact there are more efficient quantum circuit encodings: parity encoding.

[James D Whitfield, Jacob Biamonte, and Alán Aspuru-Guzik. Simulation of electronic structure hamiltonians using quantum computers. Molecular Physics, 109(5):735–750, 2011.]

Techniques

In quantum chemistry one may be interested in many different calculations: thermal rates, phase diagrams, **ground state**

There are many methods to calculate the Ground State in chemistry, such as the Adiabatic Computation (based on the Hamiltonian evolution), the Phase Estimation Algorithm (based on Hamiltonian evolution) and the **Variational Quantum Eigensolver (based just on the Hamiltonian action)**.

It was used the last one.

Variational Quantum Eigensolver




Figure: Scheme of a VQE procedure

[Nikolaj Moll, Panagiotis Barkoutsos, Lev S Bishop, Jerry M Chow, Andrew Cross, Daniel J Egger, Stefan Filipp, Andreas Fuhrer, Jay M Gambetta, Marc Ganzhorn, et al. Quantum optimization using variational algorithms on near-term quantum devices. arXiv preprint arXiv:1710.01022, 2017.]

IBM has its own quantum device, making available 14 real qubits for everyone who wants to try them.

It is also possible to use a quantum simulator with 32 qubits.

A 50 real qubits quantum computer will be available soon.



Results - H₂ molecule

| Ground State Energies | | | | |
|-----------------------|-------|-------|------------------|-----------|
| | | | ExactEigensolver | |
| | | | -1,137306035753 | |
| | | | VQE method | |
| max_trials | depth | shots | Value | Variation |
| 1 | 1 | 1 | -0,712503938173 | 4,2E-01 |
| | | 2048 | -0,102521694036 | 1,2E+00 |
| | 7 | 1 | -0,313978701260 | 8,2E-01 |
| | | 2048 | -0,593980785118 | 5,4E-01 |
| 5000 | 1 | 1 | -1,137262155450 | 4,4E-05 |
| | | 2048 | -1,137304046500 | 2,0E-06 |
| | 7 | 1 | -1,137304559058 | 2,3E-06 |
| | | 2048 | -1,137304311791 | 1,7E-06 |

Figure: The ground state energy is the lowest where a physical system can be. The energy unit used is the *Hartree*, an atomic unit of energy. The variation is the absolute error between the values obtained through Exacteigensolver and VQE method.

Results - H_2 molecule




Figure: Variation of the H_2 dissociation energy as the distance between atoms increases; parameters used in the VQE method: max_trials=350, depth=3 and shots=2048.

Results - LiH molecule

| Ground State Energies | | | | |
|-----------------------|-------|-------|-----------------|-----------|
| | | | ExactEigsolver | |
| | | | -7,88240193229 | |
| | | | VQE method | |
| max_trials | depth | shots | Value | Variation |
| 1 | 1 | 1 | -7,021299390490 | 8,6E-01 |
| | | 2048 | -7,191895551060 | 6,9E-01 |
| | 7 | 1 | -7,052667867887 | 8,3E-01 |
| | | 2048 | -6,973986243448 | 9,1E-01 |
| 5000 | 1 | 1 | -7,862023833205 | 2,0E-02 |
| | | 2048 | -7,857451742601 | 2,5E-02 |
| | 7 | 1 | -7,880613192926 | 1,8E-03 |
| | | 2048 | -7,879721385780 | 2,7E-03 |

Figure: The ground state energy is the lowest where a physical system can be. The energy unit used is the *Hartree*, an atomic unit of energy. The variation is the absolute error between the values obtained through Exacteigsolver and VQE method.

Expectations for the Future

So where do Quantum computation stands now?




Figure: Number of Qubits per molecule - IBM Research

Expectations for the Future

So where do we stand now?

Chemistry constitutes a natural field of application for quantum mechanics. We are interested in applying this methodology to study physical phenomena on the effects of physical fields over molecules.

It also seems feasible to study simplified models of biology. Quantum physics is quite relevant in many biological phenomena:

- photosynthesis, vision, smell...
- Origins of life? Consciousness?¹

¹[Schrödinger, E. (1992). What is life?: With mind and matter and autobiographical sketches. Cambridge University Press.]

Questions?

With more qubits great things are expected!

